



# The impact of infant feeding practices on the risk of developing food allergies

Wpływ sposobu żywienia niemowląt na ryzyko rozwoju alergii pokarmowej

Joanna Smorońska-Rypel<sup>1,A-F</sup>, Małgorzata Rodak<sup>1,B-C,E-F</sup>, Magdalena Kajzar<sup>1,C-F</sup>,  
Magdalena Iwan<sup>2,B-D,F</sup>, Jakub Milczarek<sup>2,A-B,E-F</sup>, Ksawery Adamiec<sup>2,B-D,F</sup>,  
Natalia Piątkowska<sup>3,B-D,F</sup>, Błażej Szymczuk<sup>2,B-D,F</sup>, Dominik Trojanowski<sup>2,B-D,F</sup>, Kamila Nitka<sup>4,B-D,F</sup>

<sup>1</sup> Department of Internal Medicine, Bonifraters Medical Center, Katowice, Poland

<sup>2</sup> Faculty of Medicine, Medical University of Silesia in Katowice, Poland

<sup>3</sup> Department of Internal Medicine, Provincial Hospital in Poznań, Poland

<sup>4</sup> Department of Internal Medicine, Łańcut Medical Center, Poland

A – Research concept and design, B – Collection and/or assembly of data, C – Data analysis and interpretation, D – Writing the article, E – Critical revision of the article, F – Final approval of the article

Smorońska-Rypel J, Rodak M, Kajzar M, Iwan M, Milczarek J, Adamiec K, Piątkowska N, Szymczuk B, Trojanowski D, Nitka K. The impact of infant feeding practices on the risk of developing food allergies. *Med Srodow*. doi: 10.26444/ms/195943

## Abstract

**Introduction.** Food allergy in infants is a major health problem in urbanised countries, the incidence of which is increasing and is a challenge for both parents and doctors. In the first months of life, a child's body is particularly sensitive to allergic reactions, therefore knowing the risk factors is crucial for the prevention and management of this health problem.

**Objective.** The aim of this study is to examine the impact of infant feeding on the development of food allergies, with particular emphasis on breastfeeding, formula milk, the method of expanding the diet, vitamin D supplementation and probiotics. The potential benefits and limitations of each method be presented.

**Brief description of the state of knowledge.** In scientific studies, there is a significant indication of the positive impact of breastfeeding on the proper development of the infant's intestinal microbiome, which is associated with a reduced risk of developing food allergies. However, the effect of formula feeding, as well as vitamin D supplementation, remains unclear and depends on many variables. Appropriate supplementation with probiotics may have a beneficial effect on the proper development of the intestinal microbiome, thereby reducing the risk of food allergies. It is not recommended to avoid or delay the introduction of allergenic foods to infants.

**Summary.** The use of appropriate infant feeding methods may reduce the risk of food allergy in infants. However, further research is needed on the effectiveness of these interventions and their optimal strategies as part of comprehensive infant care.

## Key words

breastfeeding, microbiome, probiotic, formula, food allergy in infants, expanding the diet of infants

## Streszczenie

**Wprowadzenie i cel pracy.** W krajach zurbanizowanych alergii pokarmowa u niemowląt jest poważnym problemem zdrowotnym, którego częstość występowania rośnie i który stanowi wyzwanie zarówno dla rodziców, jak i lekarzy. W pierwszych miesiącach życia dziecka jego organizm jest szczególnie wrażliwy na reakcje alergiczne, dlatego znajomość czynników ryzyka jest kluczowa dla zapobiegania alergii i zarządzania tą problematyką zdrowotną. Celem niniejszej pracy jest zbadanie wpływu sposobu karmienia niemowląt na rozwój alergii pokarmowej, ze szczególnym uwzględnieniem karmienia mlekiem kobiecym oraz mlekiem modyfikowanym, a także sposobu rozszerzania diety oraz suplementacji witaminą D i probiotykami. Przedstawia również potencjalne korzyści i ograniczenia poszczególnych metod.

**Opis stanu wiedzy.** W badaniach naukowych widoczna jest istotna zależność między pozytywnym wpływem karmienia piersią na prawidłowy rozwój mikrobiomu przewodu pokarmowego u niemowląt, co wiąże się ze zmniejszonym ryzykiem rozwoju alergii pokarmowej. Natomiast wpływ karmienia mlekiem modyfikowanym, a także suplementacji witaminą D jest niejasny i jest zależny od wielu zmiennych. Odpowiednia suplementacja probiotykami może mieć korzystny wpływ na prawidłowy rozwój mikrobiomu jelitowego, a tym samym na redukcję ryzyka wystąpienia alergii pokarmowej. Nie należy unikać ani opóźniać wprowadzania produktów w celu prewencji alergii pokarmowej u niemowląt.

**Podsumowanie.** Zastosowanie odpowiednich metod żywieniowych może zmniejszyć ryzyko wystąpienia alergii pokarmowej u niemowląt. Jednakże potrzebne są dalsze badania nad skutecznością tych działań oraz ich optymalnymi strategiami w ramach kompleksowej opieki nad niemowlętami.

## Słowa kluczowe

karmienie piersią, mikrobiom, probiotyk, mleko modyfikowane, alergii pokarmowe u niemowląt, rozszerzanie diety niemowląt

✉ Address for correspondence: Joanna Smorońska-Rypel, Department of Internal Medicine, Bonifraters Medical Center, Katowice, Poland  
E-mail: joanna.smoronska123@gmail.com

## INTRODUCTION

Food allergy in infants is a major health problem in urbanized countries, the incidence of which is increasing, leading to a challenge for both parents and doctors. Food allergy is an abnormal reaction of the immune system to contact with a potentially harmless antigen present in food. The most common type of allergic reaction is food allergy mediated by immunoglobulin E. Even the smallest amount of allergen can cause life-threatening anaphylaxis [1]. Food allergies are often confused with food intolerance, which manifests itself similarly; however, its mechanism is completely different, as it occurs without the participation of immunological mechanisms. An example of this is lactose intolerance, caused by the absence or deficiency of the enzyme lactase, necessary for the breakdown of lactose. In this case, unlike allergic reactions, small amounts of food may not cause symptoms [2].

The development of food allergy is influenced by many factors, including genetic predisposition, exposure to the allergen, immaturity of the gastrointestinal tract and severe atopic dermatitis. In the first months of life, a child's body is particularly sensitive to allergic reactions, which may lead to numerous symptoms: skin, gastrointestinal, as well as respiratory. The older the child, the lower the risk of allergies. Accurately determining the incidence of food allergy in infants is very difficult. This stems from both underestimation (studies based on data from oral challenge tests) and overestimation (data based on parents' reports of the diagnosis or occurrence of allergy symptoms) of this condition [3]. Based on numerous studies, it was estimated that the incidence of food allergy in infants, confirmed by a food challenge test, was over 10%. Although these estimates were based on food provocation tests, they were mostly performed in an open or single-blinded fashion [4]. The multicentre EuroPrevall cohort trial examined the prevalence of food allergy in European children. In the study, 12,049 newborns were enrolled, including 1,513 from Poland. Of all children, 9,336 (1,280 from Poland) were followed up at 2 years of age. In the study, the incidence of food allergy after a double-blind placebo-controlled food challenge (DBPCFC) was: for eggs 0.84% (adjusted value 1.23%) in Europe and 0.6% in Poland, for cow's milk 0.54% (adjusted value 0.74%) in Europe and 0.6% in Poland [5, 6].

Food allergy can significantly affect the quality of life, both for the child and its caregivers, requiring frequent visits to the doctor, often tedious diagnostics and the use of elimination diets, which may lead to deficiencies. The disturbing phenomenon of the rising occurrence of food allergy makes it essential for both paediatricians and parents to sufficiently understand this issue.

## OBJECTIVE

The aim of this study is to examine the impact of infant feeding on the development of food allergy, with particular emphasis on breastfeeding, formula feeding, expanding children's diet, and vitamin D and probiotic supplementation. By analyzing available scientific research and current literature, the intention is to identify differences in the risk of food allergy in infants fed using different methods, as well as to understand the mechanisms that may underlie

these relationships. Another aim is to present the potential benefits and limitations of individual feeding methods and supplementation in the context of food allergy prevention in infants, which may have significant implications for clinical practice and recommendations regarding infant feeding.

## MATERIALS AND METHOD

A review of the current literature was made in the PubMed and Google Scholar databases using the term: 'food allergy' and a combination of the term with key words such as: 'infants', 'breastfeeding', 'formula feeding', 'expanding the diet of infants', 'vitamin D', 'probiotic' and 'microbiome'. Current World Health Organization (WHO) guidelines on breastfeeding, expanding infants' diet, vitamin D, and probiotic supplementation were also reviewed. Finally, 42 original and review articles were included in the analysis.

## DESCRIPTION OF THE STATE OF KNOWLEDGE

**Breastfeeding.** The WHO recommends that infants should be exclusively breastfed for the first 6 months of life. After this period, complementary foods should be introduced and breastfeeding shall be continued until the age of 2 or even longer [7,8]. Recently, there has been a lot of pressure on the promotion of breast milk feeding due to the many beneficial effects of this method for both the child and the mother. Among other things, breastfeeding supports the development of a strong immune system which reduces the risk of acquiring many diseases, including allergies and asthma [9]. Breast milk contains essential nutrients and numerous bioactive substances, including antigen-immunoglobulin G immune complexes. It is suggested that immune complexes transferred from sensitized mothers induce tolerance in the newborn by activating its receptors and inducing antigen-specific regulatory cells. This is one of the many impressive, irreplaceable features of human milk, supporting primary prevention and subsequent proper development of the baby [10]. Breast milk also contains probiotics and prebiotics, which have beneficial effect on the development and maintenance of healthy intestinal microbiome in children. More and more research shows how crucial the proper composition of the gastrointestinal microbiota is for maintaining proper functions of the immune system. Data suggest that breastfeeding may lead to the growth of colonies of specific *Bifidobacterium* species through the prebiotic effects of human milk oligosaccharides (HMO) [11], and breastfeeding-associated bifidobacteria may support proper immune function through the production of aromatic lactic acids [12].

A study published in 2022 examined 135 healthy pregnant women and their newborns, suggested that changes in the microflora of breast milk may influence the development of food allergies. Breast milk and infant stool samples were compared at 5 time points and analyzed by microbial 16S rRNA gene sequencing. As a result, they showed that the microflora of mother's milk was more diverse and numerous in the group of infants without food allergies. A high presence of *Bifidobacterium*, *Akkermansia* and butyrate-producing bacteria was found in this group. However, the food allergy group showed a higher content of *Proteobacteria*,

*Acinetobacter* and *Pseudomonas* in mother's milk. Analysis of changes in the intestinal microflora in both groups of infants revealed compliance of changes in the number of *Bifidobacterium* with changes in the microflora of mother's milk [13]. Henrick BM et al. examined 347 stool samples taken from 157 of 208 infants born in a hospital in Stockholm, Sweden, showed that the composition of gut bacteria was highly variable at birth, but became more uniform over time. Moreover, this trial showed that the increase of *Bifidobacterium* occurred mainly in breastfed infants [14].

A study by Fujimura et al. analyzing stool samples using 16s rRNA gene sequencing in a group of 298 infants aged 1–11 months from a US birth cohort, demonstrated that the neonatal gut microbiota influences susceptibility to allergic asthma in children. happens through changes in the gut microenvironment that affect the population and function of CD4 + T cells. The group of newborns with the highest risk of developing IgE antibodies in response to allergens showed a lower relative abundance of certain bacteria (e.g. *Bifidobacterium*, *Akkermansia* and *Faecalibacterium*), and a higher relative abundance of individual fungi (*Candida* and *Rhodotorula*) in the intestinal microbiome [15].

The presented information suggests that the composition of breast milk microflora may reduce the risk of developing food allergy in infants by influencing the colonization of intestinal microflora and butyrate production. However, this topic is still under research and many factors, such as genetics, environment and the mother's diet, may also play a role in this process.

**Formula feeding.** The issue of what effect supplementing with formula milk has on the intestinal microbiome and thus on the risk of allergy should also be discussed. There is no clear evidence that formula feeding compared to breastfeeding increases the risk of food allergy. However, formula milk (although its composition is similar to mother's milk) does not contain all bioactive ingredients. Research suggests that formula-fed babies may have a different gut microbiome composition compared to exclusively breastfed babies. In the United States, 16s RNA gene sequencing was performed on stool samples from 10 healthy, exclusively breast-fed premature infants and 10 healthy formula-fed premature infants. *Propionibacterium*, *Veillonella*, and *Escherichia* were found to be abundant in the gut microbiome of breastfed infants, followed by *Staphylococcus*, *Enterococcus*, *Clostridium*, *Streptococcus*, and *Klebsiella*. However, in infants fed with formula milk, the intestines were colonized mainly by *Streptococcus* and *Klebsiella* and much less numerous *Propionibacterium* [16]. Similar results were obtained in a study on 20 Korean newborns. In the case of formula-fed infants, the microbiome may be more diverse, additionally containing numerous strains of *Escherichia coli*, *Enterococcus faecalis* and *Veillonella atypical*, and a much smaller group of *Bifidobacterium longum* [17].

In recent years, technological developments have enabled the inclusion of 2'-fucosyllactose (2'FL), the most common naturally occurring HMO, in infant formulas. Interestingly, emerging research suggests formulas enriched with HMOs have led to infants having a similar microbiome composition to that of exclusively breastfed infants [18, 19].

In the last 2 decades, access to formula milk and its consumption has increased significantly. As previously mentioned, the incidence of food allergies in infants has

increased in recent years, and one of the most common causes is cow's milk allergy (CMA) [20]. Despite the available partially hydrolyzed formula (pHF) and extensively hydrolyzed formula (eHF) used prophylactically in infants with documented family history, the problem is still growing. Amino Acid-Based Formula (AAF) is also available, which is intended for infants with severe symptoms of CMA, especially those whose symptoms persisted despite the use of eHF, or with an episode of anaphylactic shock. A recent prospective, randomized, double-blind, controlled clinical study was conducted at 20 sites in 6 countries, demonstrated the effectiveness of an elimination diet with AAF in the treatment of CMA. The trial included 169 infants with confirmed IgE-mediated CMA who were randomly assigned to 2 groups to receive AAF and AAF with a specific synbiotic blend for 12 months. Symptoms disappeared in children after 12 months of using AAF, and tolerance to cow's milk protein was achieved after 36 months of therapy in 75% of the patients. This study confirmed the effectiveness of the AAF intervention, regardless of whether it is used with or without the specific synbiotic blend, and proves that lack of exposure to cow's milk protein antigens does not slow down the development of tolerance [21].

It should be added that the European Academy of Allergy and Clinical Immunology (EAACI) guidelines suggest avoiding the administration of CMA-containing formula in the first week of life to prevent cow's milk allergy in infants and young children. However, for infants who need a breast milk substitute, there are no recommendations for or against the use of regular formula after 1 week of age to prevent food allergy [22]. Therefore, the impact of formula feeding on the intestinal ecosystem is complex and may vary depending on many factors, such as the type of formula, feeding method, family burdens, and the overall health of the child.

**Expanding the diet of infants.** Many parents are concerned about introducing new food products into their infant's diet, especially those that are most likely to cause allergies. This is often related to the fear of developing violent symptoms of allergic reactions, including life-threatening anaphylaxis.

According to the recommendations ESPGHAN (European Society for Paediatric Gastroenterology Hepatology and Nutrition), the introduction of complementary products can be started no earlier than 17 weeks of age and no later than 26 weeks of age, preferably when the infant demonstrates the developmental skills necessary to consume them [23]. However, according to WHO recommendations, expanding the diet should start from the end of 6<sup>th</sup> month of life. Until recently, there were guidelines in force indicating the exact order in which foods should be introduced. However, scientific studies conducted on large groups of children have shown that delayed introduction of potentially allergenic foods does not reduce the risk of food allergy [24]. Therefore, the current guidelines allow parents great freedom in choosing when and what food to introduce. It is recommended to introduce foods gradually, in small amounts at first, and monitor the child's reaction [25].

When examining food allergies in infants, scientists often focus on 3 main allergens: eggs, nuts and cow's milk protein, which are the most frequently reported allergy triggers in this age group [26, 27]. However, it is important to remember that any food can sensitize. Recently, several studies focused on the relationship between the moment



of introducing potentially allergenic products and the risk of allergic diseases. A meta-analysis of 5 studies (1,915 participants) conducted by Ierodiakonou et al., showed that early introduction of eggs at 4–6 months of age was correlated with a reduced risk of egg allergy. The above systematic review was presented in accordance with the PRISMA guidelines [28]. Furthermore, scientific data from two studies proved that introducing peanuts into the diet of a child aged 4–11 months was associated with a reduced risk of peanut allergy, compared to introducing them later. The Learning Early about Peanut Allergy (LEAP) Study was a randomized, controlled trial conducted in the United Kingdom that included 640 infants aged 4–11 months with severe eczema or egg allergy, then randomly assigned them to eat or avoid peanuts until age 60 months. The study showed that among infants who initially had negative results on the skin-prick test, the prevalence of peanut allergy at 60 months of age was 13.7% in the avoidance group and 1.9% in the consumption group [29].

Another randomized trial conducted in the UK by Michael R. et al. recruited 1,303 exclusively breastfed infants at 3 months of age and randomly assigned them to 2 groups: early introduction of 6 allergens or exclusive breastfeeding until 6 months of age. In a per-protocol analysis, the incidence of any food allergy was significantly lower in the early introduction group (2.4%) compared with the exclusively breastfed infant group (7.3%), as was the incidence of peanut allergy (0% vs. 5%) and egg allergy (1.4% vs. 5.5%) [30].

In general, expanding the diet in infants may increase the risk of food allergy, especially if food allergens are introduced too late. Therefore, parents should not avoid or delay introducing strong allergens into their children's diet.

**Vitamin D supplementation.** This vitamin, known mainly for its role in calcium and bone physiology, also plays an important role in the functioning of the immune system by reducing excessive inflammatory responses and promoting immune tolerance [31]. In recent years, there has been increasing interest in the possible relationship between vitamin D deficiency and the risk of food allergy in infants.

A meta-analysis conducted by Psaroulaki showed that vitamin D deficiency in mothers and infants was associated with a higher risk of food allergy, especially in the second year of life. The disease risk was 68% higher for children who did not have optimal level blood serum concentration of 25-hydroxyvitamin D [25(OH)D] [32]. The Australian randomized controlled trial examined the effects of vitamin D supplementation in early childhood on the risk of developing food allergies: it involved 195 infants with positive family history of allergic disease and adequate 25(OH)D levels at birth. Each mother supplemented vitamin D during pregnancy, reaching its serum norm in the last weeks before

delivery. The study did not show any effects of vitamin D supplementation on the risk of allergy in the first months of life. It should be taken into account that 95% of the examined infants were exclusively breastfed, therefore in the placebo group, the demand for vitamin D could be partially covered by the supply of mother's milk. This may suggest that the correct level of the vitamin in the prenatal period may be crucial in the prevention of food allergy [33].

Other studies have examined the effect of prenatal vitamin D supplementation on the risk of asthma in children. In the American randomized, double-blind trial 9268 pregnant women were assessed for eligibility and 806 had children who were included in the analysis. Children born to mothers supplementing vitamin D suffered from asthma much less often [34]. The same conclusion was reached by other American scientists who conducted randomized, double-blind, placebo-controlled trial of 806 mother-offspring pairs [35]. A similar correlation between prenatal vitamin D supplementation and the risk of food allergy was expected. A literature review from 2023 demonstrated that many research results are contradictory and no effect of vitamin D supplementation on reducing the risk of food allergy has been demonstrated. Moreover, it has been shown that both low and too-high concentrations of 25(OH)D in the prenatal period are risk factors for the occurrence of food allergy in early life [36].

Thus, although evidence indicates that 25(OH)D deficiency both prenatally and in early childhood is a risk factor for food allergy, there are no clear studies demonstrating the preventive effects of vitamin D supplementation on reducing the incidence of food allergy in infants and young children. Despite this, current recommendations state that vitamin D supplementation is necessary from the first days of life, regardless of prenatal supplementation. This is due to its proven numerous benefits for the development of the body and common deficiencies.

**Probiotic supplementation.** One of the key aspects of allergy research is the analysis of the intestinal microbiome and its impact on the functioning of the immune system. It is a complex community of microorganisms responsible not only for digestive processes but also for the immune system's functioning. The gastrointestinal microbiota influences the maturation of naive and regulatory T lymphocytes and the production of anti-inflammatory cytokines, contributing to the development of immunological tolerance and maintaining immune homeostasis [39]. Disturbing this balance, for example, through antibiotics or excessive hygiene, may increase the risk of allergies.

As previously noted, the hypothesis regarding the intestinal microbiome suggests that changes in its composition, especially in early life, may be associated with the increasing number of cases of food allergy. With the modernization of

**Table 1.** Recommended daily intake of vitamin D according to the Polish Guidelines and National Institutes of Health (NIH) [35, 36]

	Recommended Dietary Allowances for Vitamin D (Polish Guidelines)	Recommended Dietary Allowances for Vitamin D (NIH)
<b>Premature babies</b> until they reach the corrected age (according to the doctors decision)	10–20 mcg (400–800 IU)	10 mcg (400 IU)
<b>0–6 months</b>	10 mcg (400 IU)	10 mcg (400 IU)
<b>6–12 months</b>	10–15 mcg (400–600 IU)	10 mcg (400 IU)
<b>1–10 years</b>	15–25 mcg (600–1000 IU)	15 mcg (600 IU)
<b>Pregnant women</b> throughout pregnancy	50 mcg (2000 IU)	15 mcg (600 IU)

our lifestyle, reduced exposure to the natural environment, limited contact with animals, as well as excessive hygiene, changes in our microbiome may have occurred. It is believed that the higher number of allergies is due to the loss of commensal connections with parasites and bacteria – those that previously had a beneficial impact on human evolution [40, 41]. From this perspective, the use of probiotics may be a promising strategy in the prevention and treatment of allergic diseases [42].

A meta-analysis by Zhang et al. showed that the combined administration of probiotics to pregnant women and later to infants, reduced the risk of food allergy in children. However, this relationship was not found when the probiotic was used only in the prenatal period or only after delivery [43]. Australian researchers performed a double-blind, placebo-controlled randomized trial to evaluate the effectiveness of oral immunotherapy with probiotics and peanuts in children diagnosed with peanut allergy. Sixty-two patients were randomly assigned to 2 groups and underwent treatment for 18 months. At the end of the study, it was shown that 89.7% of children receiving the therapy were desensitized, compared to 7.1% of those receiving placebo. A follow-up study conducted 4 years after cessation of treatment showed that the children in the probiotic immunotherapy group were significantly more likely to be able to safely continue eating peanuts than the participants in the placebo group [44].

Currently, research confirming the effectiveness of probiotics as a preventive or therapeutic agent in the case of food allergy looks promising, but requires further efforts. The routine use of probiotics in preventing food allergy is not currently recommended.

## CONCLUSIONS

Proper infant feeding practices and supplementation are important for developing the immune system. Understanding the risk factors for infant food allergies is crucial, especially in preventing and managing this health condition in children. Education of parents about the proper way of feeding infants, including the key role of breastfeeding in the development of the intestinal microbiome, and thus the immune system, and many other beneficial properties resulting from it, is essential. However, if for some reason breastfeeding is not possible, parents should be aware of the availability of different types of formula milk and also have the opportunity to get help in choosing it. Currently, due to an abundance of information, it is very difficult for parents to find reliable materials on the proper way of feeding infants, not to mention ways of preventing allergic diseases. Education on the impact of infant feeding methods on food allergy prevention can be implemented at many levels. Effective information on the differences between breast milk and formula milk, its types, as well as the advantages and consequences of its use, that is provided by midwives during antenatal classes and postnatal visits, could influence parents' decisions. During preventive appointments paediatricians monitoring the development of infants could present individually adapted methods and instructions on expanding the diet, recommend appropriate supplementation, and could also provide a brochure with crucial information. Moreover, parents could receive reliable answers to their pressing questions. Comprehensive and organized Health Protection activities can significantly

reduce the incidence of food allergies in children and improve the quality of life of their families.

Further research is needed on methods of preventing and treating food allergies with probiotics, as well as research on formula milk to modify its composition to make it as close to mother's milk as possible. Continuous improvement of guidelines and standards of care for newborns and infants is essential for upgrading the life quality of children and their families, and is therefore required for the development of a healthy and strong society.

## REFERENCES

1. Tedner SG, Asarnej A, Thulin H, et al. Food allergy and hypersensitivity reactions in children and adults-A review. *J Intern Med.* 2022 Mar;291(3):283–302. <https://doi.org/10.1111/joim.13422>
2. Yu W, Freeland DMH, Nadeau KC. Food allergy: immune mechanisms, diagnosis and immunotherapy. *Nat Rev Immunol.* 2016 Dec;16(12):751–765. <https://doi.org/10.1038/nri.2016.111>
3. Barni S, Liccioli G, Sarti L, et al. Immunoglobulin E (IgE)-Mediated Food Allergy in Children: Epidemiology, Pathogenesis, Diagnosis, Prevention, and Management. *Medicina (Kaunas).* 2020 Mar 4;56(3):111. <https://doi.org/10.3390/medicina56030111>
4. Sampath V, Abrams EM, Adlou B, et al. Food allergy across the globe. *J Allergy Clin Immunol.* 2021 Dec;148(6):1347–1364. <https://doi.org/10.1016/j.jaci.2021.10.018>
5. Xepapadaki P, Fiocchi A, Grabenhenrich L, et al. Incidence and natural history of hen's egg allergy in the first 2 years of life-the EuroPrevall birth cohort study. *Allergy.* 2016 Mar;71(3):350–7. <https://doi.org/10.1111/all.12801>
6. Schoemaker AA, Sprikkelman AB, Grimshaw KE, et al. Incidence and natural history of challenge-proven cow's milk allergy in European children--EuroPrevall birth cohort. *Allergy.* 2015 Aug;70(8):963–72. <https://doi.org/10.1111/all.12630>
7. World Health Organization. Exclusive Breastfeeding for Six Months Best for Babies Everywhere. Available online: <https://www.who.int/news/item/15-01-2011-exclusive-breastfeeding-for-six-months-best-for-babies-everywhere> (accessed on 10 September 2024).
8. Gupta A, Suri S, Dadhich JP, et al. The World Breastfeeding Trends Initiative: Implementation of the Global Strategy for Infant and Young Child Feeding in 84 countries. *J Public Health Policy.* 2019 Mar;40(1):35–65. <https://doi.org/10.1057/s41271-018-0153-9>
9. Munblit D, Peroni DG, Boix-Amorós A, et al. Human Milk and Allergic Diseases: An Unsolved Puzzle. *Nutrients.* 2017 Aug 17;9(8):894. <https://doi.org/10.3390/nu9080894>
10. Ohsaki A, Venturelli N, Buccigrosso TM, et al. Maternal IgG immune complexes induce food allergen-specific tolerance in offspring. *J Exp Med.* 2018 Jan 2;215(1):91–113. <https://doi.org/10.1084/jem.20171163>
11. Sakanaka M, Gotoh A, Yoshida K, et al. Varied Pathways of Infant Gut-Associated Bifidobacterium to Assimilate Human Milk Oligosaccharides: Prevalence of the Gene Set and Its Correlation with Bifidobacteria-Rich Microbiota Formation. *Nutrients.* 2019 Dec 26;12(1):71. <https://doi.org/10.3390/nu12010071>
12. Laursen MF, Sakanaka M, von Burg N, et al. Bifidobacterium species associated with breastfeeding produce aromatic lactic acids in the infant gut. *Nat Microbiol.* 2021 Nov;6(11):1367–1382. <https://doi.org/10.1038/s41564-021-00970-4>
13. Wang S, Wei Y, Liu L, et al. Association Between Breastmilk Microbiota and Food Allergy in Infants. *Front Cell Infect Microbiol.* 2022 Jan 12;11:770913. <https://doi.org/10.3389/fcimb.2021.770913>
14. Henrick BM, Rodriguez L, Lakshmikanth T, et al. Bifidobacteria-mediated immune system imprinting early in life. *Cell.* 2021 Jul 22;184(15):3884–3898.e11. <https://doi.org/10.1016/j.cell.2021.05.030>
15. Fujimura KE, Sitarik AR, Havstad S, et al. Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation. *Nat Med.* 2016 Oct;22(10):1187–1191. <https://doi.org/10.1038/nm.4176>
16. Wang Z, Neupane A, Vo R, et al. Comparing Gut Microbiome in Mothers' Own Breast Milk- and Formula-Fed Moderate-Late Preterm Infants. *Front Microbiol.* 2020 May 26;11:891. <https://doi.org/10.3389/fmicb.2020.00891>
17. Lee SA, Lim JY, Kim BS, et al. Comparison of the gut microbiota profile in breast-fed and formula-fed Korean infants using pyrosequencing. *Nutr Res Pract.* 2015 Jun;9(3):242–8. <https://doi.org/10.4162/nrp.2015.9.3.242>

18. Wallingford JC, Neve Myers P, Barber CM. Effects of addition of 2-fucosyllactose to infant formula on growth and specific pathways of utilization by Bifidobacterium in healthy term infants. *Front Nutr*. 2022 Sep 23;9:961526. <https://doi.org/10.3389/fnut.2022.961526>
19. Holst AQ, Myers P, Rodríguez-García P, et al. Infant Formula Supplemented with Five Human Milk Oligosaccharides Shifts the Fecal Microbiome of Formula-Fed Infants Closer to That of Breastfed Infants. *Nutrients*. 2023 Jul 10;15(14):3087. <https://doi.org/10.3390/nu15143087>
20. Zepeda-Ortega B, Goh A, Xepapadaki P, et al. Strategies and Future Opportunities for the Prevention, Diagnosis, and Management of Cow Milk Allergy. *Front Immunol*. 2021 Jun 10;12:608372. <https://doi.org/10.3389/fimmu.2021.608372>
21. Chatchatee P, Nowak-Węgrzyn A, Lange L, et al. PRESTO study team. Tolerance development in cow's milk-allergic infants receiving amino acid-based formula: A randomized controlled trial. *J Allergy Clin Immunol*. 2022 Feb;149(2):650–658.e5. <https://doi.org/10.1016/j.jaci.2021.06.025>
22. Halken S, Muraro A, de Silva D, et al. European Academy of Allergy and Clinical Immunology Food Allergy and Anaphylaxis Guidelines Group. EAACI guideline: Preventing the development of food allergy in infants and young children (2020 update). *Pediatr Allergy Immunol*. 2021 Jul;32(5):843–858. <https://doi.org/10.1111/pai.13496>
23. Agostoni C, Decsi T, Fewtrell M, et al. ESPGHAN Committee on Nutrition. Complementary feeding: a commentary by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr*. 2008 Jan;46(1):99–110. <https://doi.org/10.1097/01.mpg.0000304464.60788.bd>
24. Ferraro V, Zanconato S, Carraro S. Timing of Food Introduction and the Risk of Food Allergy. *Nutrients*. 2019 May 21;11(5):1131. <https://doi.org/10.3390/nu11051131>
25. Szajewska H, Socha P, Horvath A, et al. Nutrition of healthy term infants. Recommendations of the Polish Society for paediatrics gastroenterology, hepatology and nutrition. *Pediatr Rev*. 2021; 50(2), 17–37. <http://dx.doi.org/10.5281/zenodo.3337389>
26. Sicherer SH, Warren CM, Dant C, et al. Food Allergy from Infancy Through Adulthood. *J Allergy Clin Immunol Pract*. 2020 Jun;8(6):1854–1864. <https://doi.org/10.1016/j.jaip.2020.02.010>
27. Peters RL, Koplin JJ, Gurrin LC, et al. HealthNuts Study. The prevalence of food allergy and other allergic diseases in early childhood in a population-based study: HealthNuts age 4-year follow-up. *J Allergy Clin Immunol*. 2017 Jul;140(1):145–153.e8. <https://doi.org/10.1016/j.jaci.2017.02.019>
28. Ierodiakonou D, Garcia-Larsen V, Logan A, et al. Timing of Allergenic Food Introduction to the Infant Diet and Risk of Allergic or Autoimmune Disease: A Systematic Review and Meta-analysis. *JAMA*. 2016 Sep 20;316(11):1181–1192. <https://doi.org/10.1001/jama.2016.12623>
29. Du Toit G, Roberts G, Sayre PH, et al. LEAP Study Team. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med*. 2015 Feb 26;372(9):803–13. <https://doi.org/10.1056/nejmoa1414850>
30. EAT Study Team. Randomized Trial of Introduction of Allergenic Foods in Breast-Fed Infants. *N Engl J Med*. 2016 May 5;374(18):1733–43. <https://doi.org/10.1056/NEJMoa1514210>
31. W Ao T, Kikuta J, Ishii M. The Effects of Vitamin D on Immune System and Inflammatory Diseases. *Biomolecules*. 2021 Nov 3;11(11):1624. <https://doi.org/10.3390/biom11111624>
32. Psaroulaki E, Katsaras GN, Samartzi P, et al. Association of food allergy in children with vitamin D insufficiency: a systematic review and meta-analysis. *Eur J Pediatr*. 2023 Apr;182(4):1533–1554. <https://doi.org/10.1007/s00431-023-04843-2>
33. Rueter K, Jones AP, Siafarikas A, et al. In “High-Risk” Infants with Sufficient Vitamin D Status at Birth, Infant Vitamin D Supplementation Had No Effect on Allergy Outcomes: A Randomized Controlled Trial. *Nutrients*. 2020 Jun 11;12(6):1747. <https://doi.org/10.3390/nu12061747>
34. Shadid IL, Brustad N, Lu M, et al. The Impact of Baseline 25-Hydroxyvitamin D Level and Gestational Age on Prenatal Vitamin D Supplementation to Prevent Offspring Asthma or Recurrent Wheezing. *Am J Clin Nutr*. 2023 Jun;117(6):1342–1352. <https://doi.org/10.1016/j.ajcnut.2023.04.019>
35. Lu M, Litonjua AA, O'Connor GT, et al. Effect of early and late prenatal vitamin D and maternal asthma status on offspring asthma or recurrent wheeze. *J Allergy Clin Immunol*. 2021 Apr;147(4):1234–1241.e3. <https://doi.org/10.1016/j.jaci.2020.06.041>
36. Feketea G, Kostara M, Bumbacea RS, et al. Vitamin D and Omega-3 (Fatty Acid) Supplementation in Pregnancy for the Primary Prevention of Food Allergy in Children-Literature Review. *Children (Basel)*. 2023 Feb 27;10(3):468. <https://doi.org/10.3390/children10030468>
37. Phudowski P, Kos-Kudła B, Walczak M, et al. Guidelines for Preventing and Treating Vitamin D Deficiency: A 2023 Update in Poland. *Nutrients*. 2023 Jan 30;15(3):695. <https://doi.org/10.3390/nu15030695>
38. National Institutes of Health. Vitamin D. Fact Sheet for Health Professionals. Available online: <https://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional/> (accessed on 10 September 2024).
39. Wang J, Zhu N, Su X, et al. Gut-Microbiota-Derived Metabolites Maintain Gut and Systemic Immune Homeostasis. *Cells*. 2023 Mar 2;12(5):793. <https://doi.org/10.3390/cells12050793>
40. Scudellari M. News Feature: Cleaning up the hygiene hypothesis. *Proc Natl Acad Sci USA*. 2017 Feb 14;114(7):1433–1436. <https://doi.org/10.1073/pnas.1700688114>
41. Poto R, Fusco W, Rinninella E, et al. The Role of Gut Microbiota and Leaky Gut in the Pathogenesis of Food Allergy. *Nutrients*. 2023 Dec 27;16(1):92. <https://doi.org/10.3390/nu16010092>
42. Cukrowska B, Ceregra A, Maciorkowska E, et al. The Effectiveness of Probiotic Lactobacillus rhamnosus and Lactobacillus casei Strains in Children with Atopic Dermatitis and Cows Milk Protein Allergy: A Multicenter, Randomized, Double Blind, Placebo Controlled Study. *Nutrients*. 2021 Apr 1;13(4):1169. <https://doi.org/10.3390/nu13041169>
43. Zhang GQ, Hu HJ, Liu CY, et al. Probiotics for Prevention of Atopy and Food Hypersensitivity in Early Childhood: A PRISMA-Compliant Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Medicine (Baltimore)*. 2016 Feb;95(8):e2562. <https://doi.org/10.1097/MD.0000000000002562>
44. Tang ML, Ponsonby AL, Orsini F, et al. Administration of a probiotic with peanut oral immunotherapy: A randomized trial. *J Allergy Clin Immunol*. 2015 Mar;135(3):737–44.e8. <https://doi.org/10.1016/j.jaci.2014.11.034>